

Brown Root Rot Resistance

Phoma sclerotoides G. Preuss ex Sacc.

F.A. Gray, C.R. Hollingsworth, R.W. Goose, C.J. Reedy and R.C. Larsen

PLANT CULTURE

Greenhouse

Container..... Stuewe and Sons, Inc., Tall One Treepots™ (top width = 10 cm, depth = 36 cm, volume = 2.83 L) (<http://www.stuewe.com>).

Media..... Pasteurized soil (1 part sand/1 part sandy loam soil).

Temp/Light..... 21C/15.6C (day/night); 12 to 16 hour day length.

No. of Plants... One plant per pot and 24 plants per rep.

Planting..... Pregerminate seed and treat with *Rhizobium meliloti* prior to planting (<http://www.nitragin.com>).

No. of Reps..... 3 to 5 replications.

Other..... Since these pots are unstable, some type of support will be required. We have used metal milk racks which hold 12 pots each.

FUNGAL ISOLATION AND INOCULUM PRODUCTION

Source..... Diseased roots. Diseased roots in plastic bags can be stored under refrigeration (4C) for 1 month.

Isolation..... The fungus can be isolated from diseased root tissue placed in water agar using standard isolation techniques. Plates must be incubated at 10C.

Production..... Culture fungus in Petri plates containing potato dextrose agar maintained at 10C. Place barley seed in a glass beaker, add water (250cc seeds/130 ml water in 600 ml beaker), cover with aluminum foil and autoclave for 60 min, allow to cool overnight and re-autoclave for an additional 60 min. When cool, transfer 2 mycelial plugs on to moistened seed and re-secure aluminum foil with parafilm. Maintain at 10C for 2 months. Following thorough colonization of barley seed, remove and spread out on clean surface and allow to air dry.

Storage..... Place barley seed inocula in plastic bags and store in freezer at -14C. Barley inocula should remain viable for up to 24 months.

INOCULATION PROCEDURE

Age of Plant.... 4 to 6 months old.

Type of Inoc... Infected barley seed containing mycelium, pycnidia and conidia.

Amount..... 2 infected barley seeds/plant. Additional inocula should increase disease development.

Procedure..... Place barley seeds against the upper tap root, approximately 2.5 cm below the soil surface. Cover inocula with a plug of sterile cotton and replace soil. Plant injury is not required.

INCUBATION

Field Plot

Culture..... After inoculation plants should be placed outside in the field in late summer to early fall. Plants are placed on top of the ground and surrounded by some type of insulation such as bales of hay to prevent roots from freezing. Root infection has been reported to occur during host dormancy from early fall to early spring. Plants should be watered as needed to maintain soil moisture.

Evaluation..... Plants are left outside through the winter months. Plants can be evaluated for root rot in late spring to early summer. The disease is inactive during the warm summer months. An additional year will result in increased disease severity. Since the suggested standard test requires plants to be placed in field plots over one-half of the year, results may vary from year to year depending on local weather conditions.

RATING PLANTS

Field Plot

Evaluating plants for severity of root rot. Partially remove plants from pots, remove soil from upper 10 cm of root, and rinse under the spigot. Rate roots for disease response on a scale of 1-5 described below:

- 1 — No disease, roots healthy.
- 2 — Slight disease, root tissues exhibit localized discoloration.
- 3 — Moderate disease, well-defined lesions on root.
- 4 — Severe root rot, plant deterioration.
- 5 — Acute root rot, plant dead.

Alfalfa populations can be characterized for reaction to Brown root rot by the percent of resistance plants (1s and 2s).

CHECK CULTIVARS

Disease Reaction	Resistant Plants (%)*			
	Expected		Acceptable Range	
	Yr 1	Yr 2	Yr 1	Yr 2
Highly Resistant Peace (Canada Dept. of Ag.)	70	50	60-80	40-60
Moderately Resistant Multi-plier (Mycogen Seeds)	20	5	10-30	0-10

*Incidence and severity of BRR should increase with increasing number of test years. Peace should be Highly Resistant for years 1 and 2, while Multi-plier should be Moderately Resistant in year 1 and Susceptible in Year 2.

SCIENTISTS WITH EXPERTISE

Name **F.A. Gray** (Plant Pathologist)
 Address College of Agriculture - Plant Sciences
 Department 3354, 1000 E. University Avenue
 University of Wyoming
 Laramie, WY 82071
 Phone (307) 766-3995
 FAX (307) 766-5549
 E-mail fagray@uwyo.edu

Name **C.R. Hollingsworth** (Plant Pathologist)
 Address Northwest Research and Outreach Center
 Department of Plant Pathology
 University of Minnesota
 Crookston, MN 56717
 Phone (218) 281-8627
 FAX (218) 281-8603
 E-mail holli030@umn.edu

Name **R.W. Goose** (Plant Breeder)
 Address College of Agriculture - Plant Sciences
 Department 3354, 1000 E. University Avenue
 University of Wyoming
 Laramie, WY 82071
 Phone (307) 766-3151
 FAX (307) 766-5549
 E-mail goose@uwyo.edu

Name **R.C. Larsen** (Plant Pathologist)
 Address USDA-ARS/PWA
 Rt. 2, Box 2953A
 Prosser, WA 99350-8694
 Phone (509) 786-3454
 FAX (509) 786-4635
 E-mail rlarsen@tricity.wsu.edu

CORRELATION TO FIELD TRIALS

A non-destructive forage yield trial, including the two BRR check cultivars, was conducted at a site in Wyoming naturally infested with *P. sclerotioides* (*P.s.*). Plant stands after three years were 86% for Peace and 16% for Multi-plier. An ongoing repeat of the previously described trial conducted at a nearby *P.s.*-infested site, has produced similar but less dramatic results. Plant stands after three years were 97% for Peace and 67% for Multi-plier. A newly established third field trial, including the two standard check cultivars and our BRR-resistant breeding population, is currently underway at a nearby site. Plants will be removed from each replicate plot after one and two years and rated for BRR severity. Results will be compared to our previous experiments using sterile inocula.

PATHOTYPES

Pathotypes of *Phoma sclerotioides* have not been reported. Studies by the authors have shown one Canadian (ATCC #56515) and 13 Wyoming isolates of *P. sclerotioides* to all be pathogenic on alfalfa.


SOURCES OF INOCULUM

Barley grain inocula of Wyoming Isolate #2, for use in the establishment of *P. sclerotioides* cultures, is available on a limited basis from the senior author. A USDA/APHIS permit is required in states where BRR has not been reported. This isolate is also available from the American Type Culture Collection (ATCC), accession number ATCC #MYA-295 (<http://www.atcc.org>).

HELPFUL INFORMATION

BRR is widespread in the Canadian Province of Saskatchewan. It also has been reported from the provinces of Alberta, British Columbia and Manitoba, as well as from the state of Alaska. In 1997, it was reported in Wyoming where it has been found to be widely distributed. It was also found in east central Idaho and has been reported from Montana. Most recently it was found in Wisconsin, Minnesota and New York. The development of a PCR based SCAR probe for *P. sclerotioides* will undoubtedly prove helpful in determining the distribution of this root pathogen in the U.S.

DISTRIBUTION AND SEVERITY OF BROWN ROOT ROT



Crossed line fill = states or provinces where extensive surveys have been conducted or multiple reports have occurred. Single line fill = states or provinces where BRR has been found but an extensive survey has not been made.

REFERENCES

- Davidson, J.G.N. 1990. Brown root rot. Compendium of Alfalfa Diseases. 2nd ed. The American Phytopathological Society, St. Paul, MN.
- Gray, F.A., C.R. Hollingsworth, C.J. Reedy, L.B. Powers and D.E. Legg. 2003. Pathogenicity of 14 isolates of *Phoma sclerotioides* pathogenic to alfalfa. Phytopathology Pacific Div. Abstracts (In Press).
- Gray, F.A., C.R. Hollingsworth, D.W. Koch and T.E. Heald. 2003. Brown root rot of alfalfa. University of Wyoming, College of Agriculture, Dept. of Plant Sciences, Laramie, WY, 82071-3354. Timely Information Series, No. 1. <http://www.uwyo.edu/plants/publications/RootRot1.htm>.

4. Gray, F.A., T.E. Heald, C.R. Hollingsworth and D.W. Koch. 1997. Brown root rot caused by *Phoma sclerotoides*, a new disease of alfalfa in the U.S. p. 22-24. In Proc. 10th West Alfalfa Improvement Conf., Davis, CA, 27-28 June 1997.
5. Hollingsworth, C.R. and F.A. Gray. 1999. First report of Brown root rot on alfalfa caused by *Phoma sclerotoides* in the continental United States. Plant Disease 83:1071.
6. Hollingsworth, C.R. 1999. Biology and management of brown root rot, *Phoma sclerotoides* of alfalfa. M.Sc. thesis, University of Wyoming, Laramie, Wyo., pp. 1-110.
7. Hollingsworth, C.R. 2002. Assessing heritability of brown root rot (*Phoma sclerotoides*) resistance and forage yield in nine alfalfa (*Medicago sativa* ssp. *sativa*) populations. Ph.D. dissertation, pp. 1-133.
8. Hollingsworth, C.R., F.A. Gray, D.W. Koch, R.W. Groose and T.E. Heald. 2003. Distribution of *Phoma sclerotoides* and incidence of brown root rot in Wyoming, U.S.A. Canadian Journal of Plant Pathology 25:215-217.
9. Hollingsworth, C.R., R.W. Groose and F.A. Gray. 2003. Comparative disease reactions of Cycle 0 and Cycle 1 alfalfa following inoculation with *Phoma sclerotoides*. Phytopathology 93:S36.
10. Larsen, R.C., C.R. Hollingsworth, G.J. Vandemark, M.A. Gritsenko and F.A. Gray. 2002. A rapid method using PCR-based SCAR markers for detection and identification of *Phoma sclerotoides*, the cause of Brown root rot disease of alfalfa. Plant Disease (86:928-932).