




Investigating Nitrogen Management in a Perennial Dual-Use Kernza-Alfalfa Intercropping System

Megan Williams
Cornell University

Kernza Intermediate Wheatgrass

Why Kernza?

- Perennial sustainability benefits
- Relatively high grain size and yield
- Consumer quality grain
- High forage biomass

Kernza Shortcomings

- Grain size is $\frac{1}{3}$ of wheat
- Grain yield decline over time



Closing the Gap Between Grasslands and Grain Agriculture (Crews, 2017)

Intercropping Kernza with Alfalfa

- ❑ Weed suppression
- ❑ Increased soil N and C in topsoil
- ❑ Refuge for beneficial insects
- ❑ Reduction in pest and disease incidence
- Yield increase
- Higher land use efficiency
- Yield stability over time



Dual-Use Cropping Systems

A dual-use system is one with the intended purpose of both harvesting grain and utilizing vegetative growth as forage for livestock.

- Improved profitability & reduced business risk (McGrath et al., 2021a).
- Bridge for both feed quality and quantity gaps at different times of the season (McGrath et al. 2021b)
- Financial risk of Kernza grain production can be reduced by managing the system for high forage yield (Hunter et al., 2020)



Experimental Objectives

Determine whether, compared to monoculture Kernza, a Kernza-alfalfa intercrop can...

- Reduce synthetic N fertilization
- Maintain or improve grain and forage yields
- Maintain or improve forage quality
- Maintain yields and quality over time
- Vary with environment



Experimental Design

4 Locations

Cornell University
(NY)

University of Wisconsin-Madison
(WI)

University of Minnesota
(MN)

The Land Institute
(KS)

2 cropping schemes

Intercrop: 5.6 kg/ha Kernza, 5.6 kg/ha Alfalfa

3 main data points

Grain Yield

Kernza Forage Biomass

3 harvests annually for 2 years

Spring pre-elongation

Summer (~m

5 Nitrogen (urea) rates

0 kg/ha

40 kg/ha

80 kg/ha

120 kg/ha

Alfalfa Variety

Shockwave BR (Fall Dormancy 4)



A photograph of a field of tall, dry grass, possibly a meadow or a field of wildflowers, with a green rectangular box in the center containing the word "Results". The background shows a line of trees and a cloudy sky.

Results


Grain Yield

N= Nitrogen Rate E= Environment C= Cropping System

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1	***	****	***	**	**	*	*
Year 2	***	***		***	***	*	*

* $P \leq 0.05$ ** $P \leq 0.01$ *** $P \leq 0.001$

- Variable response to N rate by site and cropping system
- If no differences between cropping systems, then no grain yield penalty per area in intercrop, despite half seeding rate




Total Forage Yield

N= Nitrogen Rate E= Environment C= Cropping System

Harvest	N	E	C	N:E	N:C	E:C	N:E:C	N:C	E:C	N:E:C
Year 1 Summer	***	***	***	***	***	***	***	***	15000	***
Year 1 Fall	***	***	***	***	***	***	***	***	***	***
Year 2 Summer	***	***	***	***	***	***	***	***	10000	***
Year 2 Fall	***	***	***	***	***	***	***	***	***	***

- If no response to N rate in intercrop, then opportunity to reduce N inputs and costs
- Fall Intercrop total forage yield > monoculture Kernza
- Year 1 total forage > Year 2



Cropping System

- Intercropped
- Monoculture Alfalfa
- Monoculture IWG

Harvest

- Year 1 Summer Harvest
- Year 1 Fall Harvest
- Year 2 Summer Harvest
- Year 2 Fall Harvest

Land Equivalent Ratio

Indicates intercrop yield advantage on a per-area basis


$$\text{Kernza Partial LER} = \frac{\text{Kernza yield intercrop}}{\text{Kernza yield monoculture}}$$

$$\text{Alfalfa Partial LER} = \frac{\text{Alfalfa yield intercrop}}{\text{Alfalfa yield monoculture}}$$

$$\text{LER} = \text{Kernza PLER} + \text{Alfalfa PLER}$$

LER > 1 = Yield advantage in intercrop

Replacement
Design →




LER

N= Nitrogen Rate E= Environment

Harvest	N	E	N:E
Year 1 Summer			
Year 1 Fall		***	
Year 2 Summer	**	*	
Year 2 Fall		***	

- At higher N rates, intercrop yield advantage less likely
- Intercrop confers yield advantage except at high N rate in summer harvests
- Year 2 LER can be greater than Year 1




Crude Protein (CP)

N= Nitrogen Rate E= Environment C= Cropping System

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1 Summer	**	***	***	***	**	***	***
Year 1 Fall		***	***	***	*	**	**
Year 2 Summer	*	***	***	***	**		
Year 2 Fall	***	***	***	***	**	**	
Year 2 Fall	*	***	***	***		**	**

- Intercrop CP generally did not respond to N
- Intercrop CP > monoculture Kernza CP
- Fall CP > Summer CP



Conclusions

At some locations, a farmer focused on Kernza grain yield may benefit from a dual-use Kernza-alfalfa intercropping system through....

- Higher forage biomass of higher quality in the fall
- Reduced synthetic N needs
- No significant change in grain yield at half the seeding rate of monoculture

There is more work to be done!

- Third year of results
- Alfalfa variety breeding for intercrop with Kernza (underway)
- Seeding rates in intercrop
- Economic analysis of system

Acknowledgements

Principle Investigator



Dr. Virginia Moore



Collaborators



Jake Jungers
Katherine Bohn




Brandon Schlautman
Spencer Barriball
Lee DeHaan



Valentin Picasso
Heathcliffe Riday
Ed Bures
Angad Dhariwal



National Institute of Food and Agriculture
U.S. DEPARTMENT OF AGRICULTURE

The background image shows a field of tall, dry, golden-brown grass in the foreground. In the distance, there is a green field, some trees, and a small building. The sky is overcast with grey clouds.

Questions?

Megan Williams
mlw322@cornell.edu



KS weird planting scheme

Weeds



< Perennial
weeds for
perennial
crops

Leguminous
weeds may
confound N
effect results >



Lodging May Affect Grain Yields

Monoculture

40 kg
N ha⁻¹



Intercropped




160 kg
N ha⁻¹




Weather by Site

Mean Monthly Temperature (C)



Total Monthly Precipitation (cm)



USDA Hay Quality Designation Guidelines

Alfalfa Guidelines (domestic livestock use and not more than 10% grass)

Quality	ADF	NDF	RFV	TDN-100%	TDN-90%	CP
Supreme	<27	<34	>185	>62	>55.9	>22
Premium	27-29	34-36	170-185	60.5-62	54.5-55.9	20-22
Good	29-32	36-40	150-170	58-60	52.5-54.5	18-20
Fair	32-35	40-44	130-150	56-58	50.5-52.5	16-18
Utility	>35	>44	<130	<56	<50.5	<16

Summer Intercrop

Summer
Alfalfa
Monoculture

Fall Intercrop

Fall
Monoculture
Alfalfa

Grass Hay Guidelines

Quality	Crude Protein %
Premium	Over 13
Good	9-13
Fair	5-9
Utility	Under 5




Fall monoculture kernza

Summer monoculture kernza

Land Equivalent Ratio (LER)


- Replacement Design
- Partial Land Equivalent Ratio (PLER)
- Total Land Equivalent Ratio (TLER)
- Proportional Partial Land Equivalent Ratio (PrPLER)

○ $\text{PrPLER} = \text{PLER} / \text{TLER}$




Nitrogen Rate Study: PrPLER


Year 1
Summer




Year 2
Summer



Year 1
Fall



Year 2
Fall



Kernza tended to contribute more to yield advantage in intercrop than alfalfa

Crop




Alfalfa

Kernza

Nitrogen Rate Study: Partial LER


Kernza

Harvest	N	E	N:E
Year 1 Summer		*	
Year 1 Fall		***	
Year 2 Summer			
Year 2 Fall	*	***	



Alfalfa


Harvest	N	E	N:E
Year 1 Summer		**	
Year 1 Fall		*	
Year 2 Summer	***		
Year 2 Fall	***		



Nitrogen Rate Study: Relative Feed Value (RFV) by species


Kernza

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1 Summer	***	***				*	
Year 1 Fall		***	**	**			
Year 2 Summer	*	***				***	
Year 2 Fall	**	**	*			*	



Alfalfa

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1 Summer		***				***	*
Year 1 Fall		***	***			*	
Year 2 Summer		***					
Year 2 Fall					*		




Nitrogen Rate Study: Crude Protein (CP) by species

Kernza

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1 Summer	***	***	***	***		*	
Year 1 Fall		***	***	*			
Year 2 Summer		***	**	**		*	
Year 2 Fall	***	***	***	***	*		


Relationship Between IWG CP and N-rate



Alfalfa

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1 Summer		***				**	*
Year 1 Fall		***	**		*		
Year 2 Summer		*					
Year 2 Fall		**					

Relationship Between Alfalfa CP and N-rate




Variety Study: Partial LER


Kernza

Model	G	E	G:E
Year 1 Summer		*	
Year 1 Fall		***	
Year 2 Summer			
Year 2 Fall		***	

IPLER




IPLER



Alfalfa

Model	G	E	G:E
Year 1 Summer		***	
Year 1 Fall		***	
Year 2 Summer		**	
Year 2 Fall		**	


APLER



Variety Study: Relative Feed Value (RFV) by species


Kernza

Model	G	E	G:E
Year 1 Summer		***	
Year 1 Fall		***	
Year 2 Summer		***	
Year 2 Fall			



Alfalfa


Model	G	E	C	G:E	G:C	E:C	G:E:C
Year 1 Summer		***	***			**	
Year 1 Fall		***	***			***	
Year 2 Summer		***	***			***	
Year 2 Fall		*	*				



Variety Study: Crude Protein (CP) by species


Kernza

Model	G	E	G:E
Year 1 Summer		***	
Year 1 Fall		***	
Year 2 Summer		***	
Year 2 Fall			



Alfalfa

Model	G	E	C	G:E	G:C	E:C	G:E:C
Year 1 Summer		***	*			***	
Year 1 Fall		***	***			**	
Year 2 Summer		***	***			**	
Year 2 Fall		***					



Grain dehulled weight adjustment

Analysis program: *ilastik*


Estimated dehulled sample weight = $(0.905045351733363 * \text{imaged sample weight}) - ((1.10627799960595\text{E-}05) * \text{hull fraction} * \text{hull area}) + ((4.10249606987368\text{E-}06) * \text{naked seed fraction} * \text{naked seed area}) - ((2.01262650442053\text{E-}05) * \text{Spikelet fraction} * \text{spikelet area})$

Adjusted plot grain yield = $(\text{estimated dehulled sample weight} / \text{sample weight}) * (\text{plot grain yield weight})$

Nitrogen Rate Study: Forage yield by species


Kernza

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1 Summer	***	***	***	***		**	
Year 1 Fall	**	***		**		***	
Year 2 Summer	***	***	***	***			
Year 2 Fall	***	**	**		*	***	



Alfalfa

Harvest	N	E	C	N:E	N:C	E:C	N:E:C
Year 1 Summer	*	***	***			***	
Year 1 Fall		***				*	
Year 2 Summer		***	***		***		
Year 2 Fall	**				**		





Nitrogen Effect on Forage Yields



0 kg N ha⁻¹



40 kg N ha⁻¹



80 kg N ha⁻¹



120 kg N ha⁻¹



160 kg N ha⁻¹

Intercropped alfalfa varieties across N rates



Shockwave BR 80 kg N ha-1



WL365HQ 0 kg N ha-1



WL365HQ 80 kg N ha-1



Higest 0 kg N ha-1



Higest 80 kg N ha-1

Intercrop across N rates (year 2)



40 kg N ha-1



80 kg N ha-1



120 kg N ha-1



160 kg N ha-1

Monoculture Kernza across N rates



0 kg N ha⁻¹



40 kg N ha⁻¹



80 kg N ha⁻¹



120 kg N ha⁻¹



160 kg N ha⁻¹

Monoculture vs
Intercrop
Kernza at 160
kg N ha⁻¹

Farmer trouble harvesting
grain due to lodging.... ?



Monoculture



Intercropped

Monoculture vs Intercrop Kernza at 120 kg N ha⁻¹



Monoculture



Intercropped

Monoculture vs Intercrop Kernza at 80 kg N ha⁻¹



Monoculture



Intercropped

Monoculture vs Intercrop Kernza at 40 kg N ha⁻¹



Monoculture



Intercropped