

ALFALFA STEM NEMATODE RESISTANCE

March 1995

Ditylenchus dipsaci (Kuhn) Filipjev; R. N. Peaden, G. D. Griffin, and J. L. Kugler

PLANT CULTURE

Greenhouse

Container Bench or flat deep enough to allow root development.

Media Steam pasteurized sandy loam soil mixture

Seed Prep Scarify, surface sterilize, and germinate on filter paper or seed may be directly planted and plant counts taken.

Planting Spacing 3.0 x 1.5 cm; may be direct seeded and counted prior to inoculation, particularly for selection.

Temp/Light 25°C; 12 to 16 hour daylength; supplemental winter light required.

No. of Plants 100 plants per replication.

No. of Reps 3 to 5 replications.

Other Inoculate with *Rhizobium meliloti* Dang; fertilize and spray as necessary; care in choice of insecticides is needed because some have nematicidal activity.

INOCULUM CULTURE

Source Nematodes growing on sterile callus tissue; nematodes are extracted and prepared as a water suspension.

Storage Temp 0 to 5°C.

Storage Life Maximum of 7 days in water; callus cultures can be refrigerated for several months if callus is in good condition.

Other Use a container with a large surface area to provide oxygen exchange.

INOCULATION PROCEDURE

Plant Age 2 weeks.

Type of Inoc All stages. Concentration 200 nematodes per plant.

Method Spray on using mist atomizer or if low concentration is available they may be put on with an eye dropper directly on the cotyledonary node or leaf axil; a second inoculation helps to ensure infection.

INCUBATION

Location Greenhouse flats or bench.

Culture Maintain soil moisture; encourage crown bud development by trimming top growth several times during the test period.
Age to Rate 12 weeks.

RATING

1 Resistant No swelling or distortion

2 Resistant Slight swelling but no distinct symptoms.

3 Susceptible Moderate swelling and distortion.

4 Susceptible Severe swelling and distortion.


5 Susceptible Severe necrosis or death.

CHECK CULTIVARS

	Approximate Expected Resistance (%)	Acceptable Range of Reaction (%)
Resistant		
Vernema**	60	45-70
Lahontan**	40	30-50
Lew	32	25-45
Susceptible		
Ranger**	5	0-12
Moapa 69	1	<6

Values for resistant standards include totals of 1's and 2's.

DISTRIBUTION AND SEVERITY OF ALFALFA STEM NEMATODE



- Not known to occur.
- Occurs but is not considered a problem.
- Occasionally causes significant losses on susceptible cultivars.
- Frequently causes significant losses on susceptible cultivars.

Stem nematode, *Ditylenchus dipsaci* (Kuhn) Filipjev
(Click on the map to the left for a larger version.)

SOURCE OF INOCULUM

R. N. Peaden
USDA-ARS
Rt.2 Box 2953A
Prosser, WA 99350
(509) 786-9218

SCIENTISTS WITH EXPERTISE

G. D. Griffin
USDA/ARS
Forage and Range Res. Lab.
Utah State University
Logan, UT 84322-6300
(801) 750-3073

R. N. Peaden
USDA-ARS
Rt.2 Box 2953A
Prosser, WA 99350
(509) 786-9218

J. L. Kugler
W. L. Research Inc.
21029 Road 6 SE
Warden, WA 98857-9544
(509) 349-8864

CORRELATION TO FIELD REACTION

Field reactions will be similar to greenhouse tests. However, results may be more variable or take longer to develop.

RACES

Races of *Ditylenchus dipsaci* are known to occur. The alfalfa race can parasitize and increase the mortality rate of non-host plants, but no population of the alfalfa race has been found that can reproduce on plants other than alfalfa and sanfoin.

CULTURE OPTIONS

Nematode culture for inoculation can be obtained from plants growing in the green house or the field. It may be necessary to surface sterilize before proceeding with inoculation.⁽³⁾

INOCULATION AND RANGE OF CONDITIONS

Best results are obtained in a high humidity environment. This can be obtained by covering containers with wetted cheesecloth or clear plastic germination domes. A plastic covering over the cheesecloth minimizes evaporation and creates a high humidity atmosphere. A second inoculation with 200 nematodes per plant should be made two weeks after the initial inoculation.

HELPFUL INFORMATION

Evaporation retarding agents may be useful in the nematode suspension for inoculation preparation. An environment minimizing loss of soil moisture should be maintained. If flood irrigation is used, the soil surface should be as flat and uniform as possible. Avoid uneven watering in flats. Nematodes will become concentrated in low areas.

ALTERNATIVE METHODS

Inoculate with the nematode suspension directly into the row at the time of seeding before covering the seed. This will result in some resistant seedling death when the nematodes feed at the growth point of the seedling embryo.

Mature plants can be screened by adding inoculum directly over the crown buds and covering with soil. Maintain good soil moisture.

REFERENCES

1. Elgin, J. H., Jr., B. D. Thyr, and B. J. Hartnman. 1984. Stem Nematode Resistance. Pp.30-32 In Standard tests to characterize resistance in alfalfa cultivars. USDA-ARS. Misc. Pub. No. 1434.
2. Griffin, G. D. 1984. Nematode parasites of alfalfa, cereals, and grasses. Pp. 243-321. In Plant and Insect Nematodes. W. R. Nickle, ed. New York: Marcel Dekker.
3. Krusberg, L. R., and S. Sardanelli. 1984. Technique for axenizing nematodes. J. of Nematology. 16:348